

River Eden Sustainability Partnership – Final Report

Appendix 12 – Riverfly Data

App 12 – Location of Riverfly Sites



Strathmiglo

Riverfly Data

5


Total Samples Shown

0


Trigger Level Breaches

Scores [†]



Average Score: 11









Count Distribution †



Groups



Mayfly (Ephemeridae) ‡**Total: 2 | Average: 0****Blue-Winged Olive (Ephemerellidae) ‡****Total: 150 | Average: 30****Flat-Bodied Stone Clinger (Heptageniidae) ‡****Total: 185 | Average: 37****Olives (Baetidae) ‡****Total: 505 | Average: 101****Stoneflies (Plecoptera) ‡****Total: 15 | Average: 3****Freshwater Shrimp (Gammarus) ‡****Total: 16 | Average: 3**

† Scores in this chart are calculated from the average monthly counts for each species group. The chart includes data derived from Urban and Extended Riverfly samples. Select a site using the map or menu bar to plot its trigger level alongside the scores.

† Values in this chart represent average monthly counts for each species group. The chart includes data derived from Urban and Extended Riverfly samples.

Reports powered
by Cartographer

Data collected by The
Riverfly Partnership

Data licensed OGL3

Pitlessie

Riverfly Data

6


Total Samples Shown

0


Trigger Level Breaches

Scores †



Average Score: 10









Count Distribution



Groups



Mayfly (Ephemeridae) ‡**Total: 0 | Average: 0****Blue-Winged Olive (Ephemerellidae) ‡****Total: 25 | Average: 4****Flat-Bodied Stone Clinger (Heptageniidae) ‡****Total: 261 | Average: 44****Olives (Baetidae) ‡****Total: 567 | Average: 95****Stoneflies (Plecoptera) ‡****Total: 19 | Average: 3****Freshwater Shrimp (Gammarus) ‡****Total: 133 | Average: 22**

† Scores in this chart are calculated from the average monthly counts for each species group. The chart includes data derived from Urban and Extended Riverfly samples. Select a site using the map or menu bar to plot its trigger level alongside the scores.

‡ Values in this chart represent average monthly counts for each species group. The chart includes data derived from Urban and Extended Riverfly samples.

Reports powered
by Cartographer

Data collected by The
Riverfly Partnership

Data licensed OGL3

Cow Bridge

Riverfly Data

7


Total Samples Shown

0


Trigger Level Breaches

Scores [†]



Average Score: 9









Count Distribution



Groups



Mayfly (Ephemeridae) ‡**Total: 0 | Average: 0****Blue-Winged Olive (Ephemerellidae) ‡****Total: 29 | Average: 4****Flat-Bodied Stone Clinger (Heptageniidae) ‡****Total: 23 | Average: 3****Olives (Baetidae) ‡****Total: 133 | Average: 19****Stoneflies (Plecoptera) ‡****Total: 1 | Average: 0****Freshwater Shrimp (Gammarus) ‡****Total: 4231 | Average: 604**

† Scores in this chart are calculated from the average monthly counts for each species group. The chart includes data derived from Urban and Extended Riverfly samples. Select a site using the map or menu bar to plot its trigger level alongside the scores.

† Values in this chart represent average monthly counts for each species group. The chart includes data derived from Urban and Extended Riverfly samples.

Reports powered
by Cartographer

Data collected by The
Riverfly Partnership

Data licensed OGL3

Dairsie

Riverfly Data

6


Total Samples Shown

0


Trigger Level Breaches

Scores [†]



Average Score: 9



Count Distribution †




Groups




Mayfly (Ephemeridae) ‡


Total: 0 | Average: 0

**Blue-Winged Olive (Ephemerellidae) ‡**


Total: 102 | Average: 17

**Flat-Bodied Stone Clinger (Heptageniidae) ‡**


Total: 1 | Average: 0

**Olives (Baetidae) ‡**


Total: 570 | Average: 95

**Stoneflies (Plecoptera) ‡**

Total: 0 | Average: 0

**Freshwater Shrimp (Gammarus) ‡**

Total: 6401 | Average: 1067



† Scores in this chart are calculated from the average monthly counts for each species group. The chart includes data derived from Urban and Extended Riverfly samples. Select a site using the map or menu bar to plot its trigger level alongside the scores.

‡ Values in this chart represent average monthly counts for each species group. The chart includes data derived from Urban and Extended Riverfly samples.

Reports powered
by Cartographer

Data collected by The
Riverfly Partnership

Data licensed OGL3